arXiv:1006.4301v1 [math.GR] 22 Jun 2010

Structure of Clifford Semigroups of Matrices*

Yongwen Zhu

School of Mathematics and Information Science, Yanti University, Yantai City,
P.R. China, 264005
E-mail: zyw@ytu.edu.cn

Abstract: In this paper, we characterize completely the structure of Clifford semigroups of matrices over an arbitrary field. It is shown that a semigroups of matrices of finite order is a Clifford semigroup if and only if it is isomorphic to a subdirect product of some linear (0-)groups. Then we generalize this result to a semigroup of matrices of countably infinite order and give a similar necessary and sufficient condition for it to be a Clifford semigroup.

Key Words: matrix semigroups; Clifford semigroups; subdirect product.

MSC 2000: 20M25

1 Introduction and preliminaries

Throughout the paper, F is an arbitrary field, n is a positive integer. Let $M_n(F)$ and $GL(n, F)$ be the set of all matrices and that of all invertible matrices of $n \times n$ over F, respectively. A *matrix semigroup* is defined as a semigroup whose set is a

---

*Research supported by National Natural Foundation of China, grant 10571005.

subset of $M_n(F)$ and whose composition is the usual multiplication of matrices, see [1], [2], [3], [4], [5] and [6]. If a matrix semigroup is a group, then we call it a *matrix group*. For example, $M_n(F)$ and $GL(n, F)$ are both matrix semigroups, and the latter is even a matrix group. Evidently, any matrix semigroup must be a subsemigroup of $M_n(F)$. By a *linear group*, we mean a subgroup of $GL(n, F)$. Note that a matrix group must be isomorphic to some linear group. For instance, $\{\begin{pmatrix} x & 0 \\ 0 & 0 \end{pmatrix} : x \in F \setminus \{0\}\}$ is a group of matrices, which is isomorphic to $GL(1, F)$, the linear group of dimension 1.

Inverse semigroups are an important class of semigroups, and Clifford semigroups are an important class of inverse semigroups, see [7], [8], [9] and [10]. In fact, a *Clifford semigroup* may be defined as a completely regular inverse semigroup. We have studied the inverse semigroups of matrices in [2]. In this paper, we shall characterize completely the structure of a Clifford semigroup of matrices. Our main result of this paper is Theorem 2.1, which gives a sufficient and necessary condition for a semigroup of matrices of finite order to be a Clifford semigroup. This theorem should be generalized to those of matrices of infinite order in the last section of this paper. As a direct corollary of our main theorem, Corollary 2.5 improves the main results of [10].

For proving our main theorem in the next section, we present now some lemmas for preliminaries.

The first lemma comes from [2, Corollary 3.5].

Lemma 1.1 *If S is a Clifford semigroup of some $n \times n$ matrices with a matrix of* *rank r, then there exists an invertible matrix P so that for all $a \in S$, there exist an $r \times r$ matrix U and an $(n-r)\times(n-r)$ matrix Z such that* $P^{-1}aP = \begin{pmatrix} U & 0 \\ 0 & Z \end{pmatrix}$.

We call a matrix $a \in M_n(F)$ *fullranked* if the rank of a is n, that is $r(a) = n$. We call a semigroup $S \subseteq M_n(F)$ *fullranked* if every element of S is fullranked. If S contains the zero matrix 0 and every nonzero matrix of S is fullranked, then we say S is *0-fullranked.* If G is a group, then $G^0 = G \cup \{0\}$ is a semigroup. We call a segmiroup formed in this way a 0-*group.* Now we can prove our second lemma.

Lemma 1.2 *A (0-)fullranked regular semigroup of matrices is a (0-)group.*

Proof. Suppose that $S \subseteq M_n(F)$ is fullranked regular semigroup, that is $S \subseteq GL_n(F)$. For any nonzero element $a \in S$, a is actually an invertible matrix. So $aba = a$ implies that $ab = 1$, the identity matrix. Therefore, S is a linear group.

Similarly, one can prove that if $S \subseteq M_n(F)$ is 0-fullranked regular semigroup, then S is a 0-group. So the proof is complete

If $S \subseteq M_n(F)$ is a semigroup and $r(a) = r(b)$ for any $a, b \in S$, then we call S *monoranked.* If $S \subseteq M_n(F)$ is a semigroup containing the zero matrix 0 and $r(a) = r(b)$ for any $a, b \in S \setminus \{0\}$, then we call S *0-monoranked.* For Green's relations $\mathcal{H}$,$\mathcal{R}$,$\mathcal{L}$,$\mathcal{D}$ and $\mathcal{J}$, we refer to [7], [8] and [9]. Then we can show the next statement, the last lemma of this section.

Lemma 1.3 *A (0-)monoranked Clifford semigroup of matrices is a (0-)group.*

Proof. Suppose that $S \subseteq M_n(F)$ is 0-monoranked Clifford semigroup. Then for any $a, b \in S \setminus \{0\}$, $r(a) = r(b)$. In light of [1, Lemma 2.1], $a\mathcal{J}b$. From

[7, Theorem 4.2.1], it follows that $\mathcal{J} = \mathcal{H}$ for Clifford semigroup$S$. Thus $a\mathcal{H}b$. Therefore, $S \setminus \{0\}$ is a group and then S is a 0-group.

Similarly, one can deduce that S is a group provided that $S \subseteq M_n(F)$ is monoranked Clifford semigroup. So the proof is complete

2 Structure of Clifford semigroups of matrices of finite order

Suppose that $S \subseteq M_n(F)$ is a semigroup and there exists a matrix $P \in GL(n, F)$ such that for every $a \in S$, $P^{-1}aP = diag\{a_1, \cdots, a_t, 0, \cdots, 0\}$ with $a_k \in M_{n_k}(F)$ for $k = 1, \cdots, t$. For any $k \leq t$, let $S_k = \{a_k | a \in S\}$. Then $S_k \subseteq M_{n_k}(F)$ is a semigroup. We denote $S \sim [S_1, \cdots, S_k]$. Now we can prove our main theorem of this paper, which characterizes completely the Clifford semigroups of $n \times n$ matrices, where n is finite.

Theorem 2.1 *Let F be a field and $S \subseteq M_n(F)$ a semigroup. Then S is a Clifford semigroup if and only if S is isomorphic to a subdirect product of some linear (0-)groups.*

Proof. Sufficiency. If S is isomorphic to a direct product of some (0-)groups, then S is obviously a Clifford semigroup because every (0-)group is evidently a Clifford semigroup.

Necessity. Assume that $S \subseteq M_n(F)$ is a Clifford semigroup. If there is a matrix $a \in S$ with $0 < r(a) < n$, then by Lemma 1.1, we have $S \sim [S_1, S_2]$. The same argument can be used to S_1 and S_2 respectively. So by using repeatedly Lemma 1.1, we should assume that $S \sim [S_1, \cdots, S_t]$ with each S_i (0-)fullranked. It is easy to see that S is Clifford semigroup if and only if $S_1, \cdots, S_t$ are all Clifford semigroups. Thus $S_1, \cdots, S_t$ are all (0-)fullranked regular semigroups of matrices. In light of Lemma 1.2, each S_i is a group or a 0-group. Therefore, S is isomorphic to the subdirect product of linear (0-)groups $S_1, \cdots, S_t$ and the proof is complete.

The next corollary is a direct consequence of the last theorem, which appears also as [2, Lemma 4.1].

Corollary 2.2 *A Clifford semigroup of 2×2 matrices over a field F is commutative if it contains a matrix of rank 1.*

Corollary 2.3 *If $S \subseteq M_3(F)$ is a Clifford semigroup which contains at least one singular matrix, then every matrix of S has at least one eigenvalue in F.*

Corollary 2.4 *If $S \subseteq M_n(F)$ is a Clifford semigroup and every matrix of S is of rank 1, then S is a commutative group and all eigenvalues of every matrix of S are in F.*

Proof. Since S is monoraked, S is a group by Lemma 1.3. According to Theorem 2.1, one can easily complete the proof.

The next corollary is an immediate consequence of Theorem 2.1.

Corollary 2.5 *Let F be a field and $S \subseteq M_n(F)$ a semigroup. Then S is a maximal Clifford subsemigroup of $M_n(F)$ if and only if S is isomorphic to the $0-$group $GL_n(F) \bigcup \{0\}$.*

Note that the last corollary generalizes Theorem 7 of [10], which stated the same result only under the condition that F is a finite field.

3 Structure of Clifford semigroups of matrices of countably infinite order

Given a finite positive integer n, we define a function $\phi_n : M_n(F) \to M_{n+1}(F)$ as follows: $\phi_n(a) = \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}$ for any $a \in M_n(F)$. It is clear that ϕ_n is a homomorphism of semigroups. Let $M_{\mathcal{X}_0}(F) = \overrightarrow{\lim}(M_n(F), \phi_n)$, the direct limit of the system $(M_n(F), \phi_n)$. Then $M_{\mathcal{X}_0}(F)$ is a semigroup, called the *full matrix semigroup of countably infinite order.*

For any $s \in M_{\mathcal{X}_0}(F)$, we can view s naturally as a matrix with the size $\mathcal{X}_0 \times \mathcal{X}_0$.

If $S \subseteq M_{\mathcal{X}_0}(F)$ is a semigroup, then we call S is *matrix semigroup of countably infinite order.* Now, we can generalize our main result presented in last section to semigroups of matrices of countably infinite order and obtain the following theorem.

Theorem 3.1 *Let F be a field and $S \subseteq M_{\mathcal{X}_0}(F)$ a semigroup. Then S is a Clifford semigroup if and only if S is isomorphic to a subdirect product of some linear (0-)groups of finite order.*

This theorem can be proved by the same method as used in the proof of Theorem 2.1, but the Lemma 1.1 should be replaced by the next lemma.

Lemma 3.2 *If* $S \subseteq M_{\mathcal{X}_0}(F)$ *is a Clifford semigroup with a matrix of rank* r*, then there exists an invertible matrix* P *of countably infinite order so that for all* $a \in S$*, there exist an* $r \times r$ *matrix* U *and a matrix* Z *of countably infinite such that* $P^{-1}aP = \begin{pmatrix} U & 0 \\ 0 & Z \end{pmatrix}$.

Proof. Since S is a Clifford semigroup, S is an inverse semigroup. We denote its idempotents by E. If S contains a matrix b of rank r, then $bb^{-1} = e \in E$. It is clear that $r(e) = r(bb^{-1}) = r(b) = r$. By the definition of $M_{\mathcal{X}_0}(F)$, there exists a positive integer n such that $e = \begin{pmatrix} f & 0 \\ 0 & 0 \end{pmatrix}$ with $f \in M_n(F)$. It follows that f is an idempotent from that e is an idempotent. Evidently, $r(f) = r(e) = r$. Thus there exists $P_0 \in GL(n, F)$ so that $P_0^{-1} f P_0 = \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}$. Let $P = \begin{pmatrix} P_0 & 0 \\ 0 & I \end{pmatrix}$ where I is a matrix of countably infinite order all elements of which are 0 but the diagonals are 1s. Then P is an invertible matrix of countably infinite order, and $P^{-1}eP = \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}$.

For any $a \in S$, set $P^{-1}aP = c$. Since $P^{-1}SP$ is isomorphic to S and S is a Clifford semigroup, $P^{-1}SP$ is also a Clifford semigroup. In light of [7, Theorem 4.2.1], in a Clifford semigroup, any element and any idempotent are commutative, so $ce = ec$. Assume $c = \begin{pmatrix} U & X \\ Y & Z \end{pmatrix}$ with $U \in M_r(F)$. Then $\begin{pmatrix} U & X \\ Y & Z \end{pmatrix}\begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}\begin{pmatrix} U & X \\ Y & Z \end{pmatrix}$, that is $\begin{pmatrix} U & 0 \\ Y & 0 \end{pmatrix} = \begin{pmatrix} U & X \\ 0 & 0 \end{pmatrix}$. It follows that $X = 0$ and $Y = 0$. So $P^{-1}aP = c = \begin{pmatrix} U & 0 \\ 0 & Z \end{pmatrix}$ as desired.

As an immediate consequence of Theorem 3.1, the following corollary is similar to the Corollary 2.4.

Corollary 3.3 *If $S \subseteq M_{\mathcal{X}_0}(F)$ is a Clifford semigroup and every matrix of S is of rank 1, then S is a commutative group and all matrices of S are diagonalizable simultaneously.*

AcknowledgementThe author would like to express his gratitude to Professor Wang Efang.

References

[1] Okniński J. *Semigroups of Matrices* [M]. Singapore: World Scientific, 1998.

[2] Zhu, Yongwen. Inverse semigroups of matrices [J]. *J. of Mathematical Research and Exposition*, 2008, **28**(3): 549-557.

[3] Zhu, Yongwen. Compact abelian matrix semigroups [J]. *J. of Mathematical Research and Exposition*, 2007, **27**(3): 601-604. (in Chinese)

[4] Zhu, Yongwen. Regular matrix semigroups [J]. *Advances in Mathematics*, 2009, **38**(1): 75-78. (in Chinese)

[5] Zhu, Yongwen. Completely simple matrix semigroups [J]. *Advances of Mathematics*, 2007, **36**(1): 76-80. (in Chinese)

[6] Zhu, Yongwen. Reducibility of Commutative Semigroups of Matrices [J]. *Acta Mathematica SinicaChinese Series*, 2010, **53**(4): (in press). (in Chinese)

[7] Howie, J. M. *Fundamentals of Semigroup Theory* [M]. Oxford: Clarendon Press, 1995.

[8] Petrich M. *Inverse Semigroups*[M]. New York: John Wiley and Sons, Inc., 1984.

[9] Zhu, Yongwen. The fundamental properties of tiling semigroups [J]. J. of Algebra, 2002, **252**: 195-204.

[10] Lee, Edmond W. H. Maximal Clifford semigroups of matrices. [J]. Sarajevo J. of Mathematics, 2006, **15**(2): 147-152.